

A NEW SPIN ON GALACTIC DUST

ANGÉLICA DE OLIVEIRA-COSTA^a, MAX TEGMARK^a, DOUGLAS P. FINKBEINER^b,
 R.D. DAVIES^c, CARLOS M. GUTIERREZ^d, L.M. HAFNER^e, ALED W. JONES^f, A.N. LASENBY^f,
 R. REBOLO^d, RON J. REYNOLDS^e, S.L. TUFTE^g AND R.A. WATSON^{c,d}

^aDepartment of Physics & Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA

^bDepartment of Astronomy, Princeton University, Princeton, NJ 08544, USA

^cUniversity of Manchester, Nuffield Radio Astronomy Laboratories, Jodrell Bank, Cheshire, SK11 9DL, UK

^dInstituto de Astrofísica de Canarias, 38200 La Laguna, Tenerife, Spain

^eAstronomy Department, University of Wisconsin, Madison, WI 53706, USA

^fMullard Radio Astronomy Observatory, Cavendish Laboratory, Cambridge, CB3 0HE, UK

^gPhysics Department, Lewis and Clark College, Portland, OR 97219, USA

February 1, 2008; ApJ, in press

ABSTRACT

We present a new puzzle involving Galactic microwave emission and attempt to resolve it. On one hand, a cross-correlation analysis of the WHAM H α map with the Tenerife 10 and 15 GHz maps shows that the well-known DIRBE correlated microwave emission cannot be dominated by free-free emission. On the other hand, recent high resolution observations in the 8–10 GHz range with the Green Bank 140 ft telescope by Finkbeiner *et al.* failed to find the corresponding 8σ signal that would be expected in the simplest spinning dust models. So what physical mechanism is causing this ubiquitous dust-correlated emission? We argue for a model predicting that spinning dust is the culprit after all, but that the corresponding small grains are well correlated with the larger grains seen at $100\mu\text{m}$ only on large angular scales. In support of this grain segregation model, we find the best spinning dust template to involve higher frequency maps in the range 12–60 μm , where emission from transiently heated small grains is important. Upcoming CMB experiments such as ground-based interferometers, MAP and Planck LFI with high resolution at low frequencies should allow a definitive test of this model.

Subject headings: cosmic microwave background – diffuse radiation – radiation mechanisms: thermal and non-thermal – methods: data analysis

1. INTRODUCTION

Understanding the physical origin of Galactic microwave emission is interesting for two reasons: to determine the fundamental properties of the Galactic components, and to refine the modeling of foreground emission in Cosmic Microwave Background (CMB) experiments. There are three Galactic foregrounds currently identified: synchrotron radiation, free-free emission and thermal (vibrational) emission from dust grains. In the last few years, however, it has become clear that a fourth component exists. This component, which we nickname “Foreground X”, is spatially correlated with $100\mu\text{m}$ dust emission but with a spectrum rising towards lower frequencies in a manner that is incompatible with thermal dust emission.

This fourth component was first discovered in the COBE DMR data by Kogut *et al.* (1996a; 1996b), who tentatively identified it as free-free emission. It has since been detected in the data sets from Saskatoon (de Oliveira-Costa *et al.* 1997), OVRO (Leitch *et al.* 1997), the 19 GHz survey (de Oliveira-Costa *et al.* 1998) and Tenerife (de Oliveira-Costa *et al.* 1999, hereafter dOC99; Mukherjee *et al.* 2000). Draine and Lazarian (1998, hereafter DL98) argued against the free-free hypothesis on energetic grounds, and suggested that Foreground X was caused by dust after all, but via rotational rather than vibrational emission. The fact that the spectrum of Foreground X is observed to flatten towards lower frequencies and perhaps even turn over around 15 GHz (dOC99) agrees well with the spin-

ning dust predictions¹. Although a consensus has still not been reached on this point (Mukherjee *et al.* 2000), the case for spinning dust was beginning to look quite solid until recently.

New observations done by Finkbeiner *et al.* (2001, hereafter F01) have reopened the question about the existence and nature of Foreground X. Using the Green Bank 140 ft telescope, these authors observed 10 IRAS dust filaments at arcminute (FWHM $\sim 6'$) scales in the frequencies 5, 8 and 10 GHz². Although the Draine and Lazarian spinning dust model normalized to the Tenerife observations suggested that several filaments should be detected at the 8σ level, F01 observations shows that only two of these 10 regions are correlated with the $100\mu\text{m}$ dust map, and only one of these two detections is compatible with the Tenerife results. Moreover, their frequency spectra is consistent with spinning dust model of DL98 and inconsistent with free-free emission alone.

The goal of this *Letter* is to perform new cross-correlation tests and to present a model that reconciles these apparently contradictory results. Since the recently completed map from the Wisconsin H-Alpha Mapper (WHAM) should trace free-free emission, it offers a powerful way to test the free-free explanation of Foreground X. A recent cross-correlation analysis of WHAM with QMAP (de Oliveira-

¹ Adding information to this puzzle, correlations between dust and $H\alpha$ maps are marginal (McCullough 1997; Kogut 1997).

² The selected regions were carefully chosen to be cold neutral clouds or HII regions at low density because in both objects the free-free emission is expected to be subdominant.




FIG. 1.— The frequency dependence of rms foreground fluctuations is shown for four different Galactic latitude slices. Squares show DIRBE-correlated emission (Foreground X), triangles Haslam-correlated emission (synchrotron), and circles $H\alpha$ -correlated emission (free-free) for four Galactic latitude slices ($0^\circ - 10^\circ$, $10^\circ - 20^\circ$, $20^\circ - 30^\circ$ and $30^\circ - 40^\circ$). The free-free curve is seen to lie below Foreground X in all cases, typically by about an order of magnitude. Foreground X therefore cannot be dominated by free-free emission. The corresponding curves represent the best fit models. The 10 and 15 GHz points are from the Tenerife correlations, the 31.5, 53 and 90 GHz points are from the DMR correlations. Upper limits are $2-\sigma$.

Costa *et al.* 2000, hereafter dOC00) failed to reveal a significant correlation, but this is hardly surprising since QMAP failed to detect significant contributions of Foreground X as well. Below we will start out in §2 by cross-correlating WHAM with the Tenerife data, where Foreground X was previously detected at the 4.5σ level (dOC99). We will find that free-free emission is detected at levels far too low to explain Foreground X. In §3 we therefore proceed to propose an alternative model for what is going on.

2. CONSTRAINTS FROM THE CROSS-CORRELATION

2.1. Cross-correlation method

In this section, we quantify the foreground contamination in the Tenerife data by cross-correlating it with a number of Galactic emission templates. This is done with the multi-component fit method, described in detail in dOC99. This method models the vector of Tenerife observations $\mathbf{y}$ as a sum

$$\mathbf{y} = \mathbf{X}\mathbf{a} + \mathbf{x}_{CMB} + \mathbf{n}, \quad (1)$$

where $\mathbf{X}$ is a matrix whose rows contain the various foreground templates convolved with the Tenerife triple-beam (*i.e.*, $\mathbf{X}_{ij}$ would be the i^{th} observation if the sky had looked like the j^{th} foreground template), and $\mathbf{a}$ is a vector of size

M that gives the levels at which these foreground templates are present in the Tenerife data.

The estimate of $\mathbf{a}$ is computed by minimizing $\chi^2 \equiv (\mathbf{y} - \mathbf{X}\mathbf{a})^T \mathbf{C}^{-1} (\mathbf{y} - \mathbf{X}\mathbf{a})$, and by modeling the covariance matrix $\mathbf{C}$ including both the experimental noise $\mathbf{n}$ and CMB sample variance in the CMB signal $\mathbf{x}_{CMB}$. As in dOC99, our quoted error bars include the effects of both noise and chance alignments between the CMB and the various template maps. The minimum-variance estimate of $\mathbf{a}$ is

$$\hat{\mathbf{a}} = [\mathbf{X}^T \mathbf{C}^{-1} \mathbf{X}]^{-1} \mathbf{X}^T \mathbf{C}^{-1} \mathbf{y} \quad (2)$$

with covariance matrix

$$\boldsymbol{\Sigma} \equiv \langle \hat{\mathbf{a}} \hat{\mathbf{a}}^T \rangle - \langle \hat{\mathbf{a}} \rangle \langle \hat{\mathbf{a}}^T \rangle = [\mathbf{X}^T \mathbf{C}^{-1} \mathbf{X}]^{-1}. \quad (3)$$

2.2. Data used

We use the latest version of the Tenerife data (Mukherjee *et al.* 2000), which has more accurate offset removal than that used in dOC99. To obtain accurate estimates of foreground levels, it is important that the analysis includes all relevant emission components and the correlations between them. As a synchrotron template, we use the 408 MHz survey (Haslam *et al.* 1982, hereafter H82), and the 1420 MHz survey (Reich 1982; Reich and Reich 1986) for cross-checks. As a template for both vibrational and

spinning dust emission, we use the Diffuse Infrared Background Experiment (DIRBE) sky maps at wavelengths from 12 to 240 μm (Boggess *et al.* 1992). Finally, as a tracer of free-free emission, we use the Wisconsin H-Alpha Mapper (WHAM) survey (Haffner *et al.* 1999). The extent of point source contamination in the Tenerife data was discussed and estimated in Gutiérrez *et al.* (2000), and will therefore not be addressed in this *Letter*. In practice, we just remove the estimated point source contribution before calculating the correlations.

2.3. Cross-correlation results

Cross-correlation results are presented in Table 1 and Figure 1. All fits are done jointly for 3 templates (100 μm – H82 – WHAM), and statistically significant ($> 2\sigma$) correlations listed in Table 1 are in boldface. Since the fluctuation levels depend strongly on Galactic latitude, we perform our analysis for six different latitude slices of roughly equal area (Figure 1) as well as for three different latitude cuts (Table 1): 20°, 30° and the *Tenerife cut* (which consists of data inside the region $160^\circ < \text{RA} < 250^\circ$, corresponding to Galactic latitudes $|b| \gtrsim 40^\circ$). For definiteness, we use the DIRBE 100 μm channel when placing all limits shown in this subsection since it is the least noisy of the DIRBE channels, and the Haslam map since it is the

synchrotron template at lowest frequency.

Figure 1 shows the corresponding fluctuations in antenna temperature in the Tenerife data ($\delta T = \hat{\mathbf{a}}\sigma_{Gal}$, where σ_{Gal} is the standard deviation in the template map). The Haslam and DIRBE correlations are seen to be consistent with those from dOC99. Synchrotron emission (triangles) generally dominates the rms foreground fluctuations at 10 GHz. At 15 GHz, on the other hand, Foreground X (squares) is seen to dominate except in the Galactic plane itself. The key new result here is the inclusion of the WHAM data, showing that free-free emission (circles) is about an order of magnitude below Foreground X over the entire range of frequencies and latitudes where it is detected. This means that Foreground X cannot be explained as free-free emission. The corresponding values of a_i (in units of $\mu\text{K/R}$) are consistent with gas at 8000K (Bennett *et al.* 1992) for latitudes $|b| = 0^\circ - 10^\circ$ and frequencies ≥ 30 GHz, with substantial scatter elsewhere³.

As in de Oliveira-Costa *et al.* (1997; 1998), dOC99 and dOC00, the cross-correlation software was tested by analyzing constrained realizations of CMB and Tenerife instrument noise. From 1000 realizations, we recovered unbiased estimates $\hat{\mathbf{a}}$ with a variance in excellent agreement with equation (3). As an additional test, we computed $\chi^2 \equiv (\mathbf{y} - \mathbf{X}\mathbf{a})^T \mathbf{C}^{-1}(\mathbf{y} - \mathbf{X}\mathbf{a})$ and obtained $\chi^2/N \approx 1$ in almost all cases.

We also performed a joint fit retaining the 100 μm , Haslam and WHAM templates in the Tenerife observing region, but replacing the Tenerife data by COBE DMR data at 31.5, 53 and 90 GHz. These results are also plotted in Figure 1 by multiplying the measured coefficients $\hat{\mathbf{a}}$ by template rms σ_{Gal} corresponding to the Tenerife triple beam, and show that our conclusions extend to higher fre-

³ Although the cross-correlation technique can accurately determine the dominant components, the detailed results for strongly subdominant components must be taken with a grain of salt. This is because any complications with the dominant components (*e.g.*, slight spatial variations in their frequency dependence) which are not included in the model of equation (1) will act as excess noise on the remaining components. Although the analysis clearly demonstrates that free-free emission is subdominant, the formal error bars on this component are therefore likely to be smaller than the true errors.

Table 1 – Correlations for 10 and 15 GHz data.

$b \ \& \ \nu$	Map ^(a)	$\hat{\mathbf{a}} \pm \delta\hat{\mathbf{a}}^{(b)}$	$\frac{\hat{\mathbf{a}}}{\sigma_{Gal}}$	σ_{Gal}	$\delta T \ [\mu\text{K}]^{(c)}$
$ b > 20^\circ$ 10 GHz	100 μm	71.4 ± 14.2	5.0	0.7	49.9 ± 9.9
	H82	71.3 ± 6.3	11.3	1.0	71.3 ± 6.3
	WHAM	-4.7 ± 19.9	-0.2	0.4	-1.9 ± 8.0
15 GHz	100 μm	51.4 ± 8.0	6.4	0.7	35.9 ± 5.6
	H82	4.1 ± 4.1	1.0	1.0	4.1 ± 4.1
	WHAM	-13.6 ± 10.8	-1.3	0.4	-5.4 ± 4.3
$ b > 30^\circ$ 10 GHz	100 μm	-3.5 ± 40.0	-0.1	0.2	-0.7 ± 8.0
	H82	5.8 ± 9.4	0.6	0.9	5.2 ± 8.5
	WHAM	39.3 ± 75.2	0.5	0.1	3.9 ± 7.5
15 GHz	100 μm	53.7 ± 26.3	2.0	0.2	10.7 ± 5.3
	H82	-4.5 ± 6.0	-0.8	0.9	-4.0 ± 5.4
	WHAM	-38.1 ± 43.6	-0.9	0.1	-3.8 ± 4.3
$ b \gtrsim 40^\circ$ 10 GHz	100 μm	65.9 ± 60.5	1.1	0.2	13.2 ± 12.1
	H82	-4.8 ± 12.3	-0.4	0.8	-3.8 ± 9.8
	WHAM	19.7 ± 107.0	0.2	0.1	1.9 ± 10.7
15 GHz	100 μm	27.0 ± 43.3	0.6	0.2	5.4 ± 8.7
	H82	-3.6 ± 7.5	-0.5	0.8	-2.8 ± 6.0
	WHAM	-55.0 ± 58.2	-0.9	0.1	-5.5 ± 5.8
$ b > 20^\circ$ 10 GHz	Fink12 μm	777.3 ± 273.8	2.8	0.03	23.3 ± 8.2
	H82	73.1 ± 6.3	11.6	1.0	73.1 ± 6.3
	WHAM	5.0 ± 19.8	0.3	0.4	2.0 ± 7.9
15 GHz	Fink12 μm	763.2 ± 169.7	4.5	0.03	22.9 ± 5.1
	H82	5.2 ± 4.1	1.3	1.0	5.2 ± 4.1
	WHAM	-14.0 ± 10.9	-1.3	0.4	-5.6 ± 4.4
$ b > 20^\circ$ 10 GHz	Combo	1.0 ± 0.2	5.0	58.1	58.1 ± 11.7
	H82	68.5 ± 6.4	10.7	1.0	68.5 ± 6.4
	WHAM	-15.2 ± 20.2	-0.8	0.4	-6.1 ± 8.1
15 GHz	Combo	1.0 ± 0.2	5.0	32.4	32.4 ± 6.5
	H82	4.5 ± 4.1	1.1	1.0	4.5 ± 4.1
	WHAM	-13.9 ± 10.8	-1.3	0.4	-5.6 ± 4.3

(a) The DIRBE, Haslam and WHAM correlations listed above correspond to joint 100 μm –H82– H_α fit.

(b) $\hat{\mathbf{a}}$ has units $\mu\text{K} (\text{MJy/sr})^{-1}$ for the 100 μm , $\mu\text{K/K}$ for the Haslam and $\mu\text{K/R}$ for the WHAM template.

(c) $\delta T \equiv (\hat{\mathbf{a}} \pm \delta\hat{\mathbf{a}})\sigma_{Gal}$.

Table 2 – 100 μm – H_α correlations.

Authors ^(a)	b	$\hat{\mathbf{a}} \pm \delta\hat{\mathbf{a}}^{(b)}$	$\frac{\hat{\mathbf{a}}}{\sigma_{Gal}}$
McCullough (1997)	$b = -65^\circ$	0.79 ± 0.24	3.3
Kogut (1997)	$b = -21^\circ$	0.85 ± 0.44	1.9
	$b = +27^\circ$	0.34 ± 0.33	1.0
dOC00	$b = +27^\circ$	0.25 ± 0.19	1.3
This work ^(c)	$ b > 20^\circ$	0.18 ± 0.07	2.6
	$ b > 30^\circ$	0.17 ± 0.04	4.3
	<i>Tenerife Cut</i>	0.04 ± 0.03	1.3
	$00^\circ \lesssim b \lesssim 10^\circ$	0.004 ± 0.100	0.04
	$10^\circ \lesssim b \lesssim 20^\circ$	0.04 ± 0.17	0.2
	$20^\circ \lesssim b \lesssim 30^\circ$	0.16 ± 0.15	1.1
	$30^\circ \lesssim b \lesssim 42^\circ$	0.16 ± 0.03	5.3
	$42^\circ \lesssim b \lesssim 56^\circ$	0.14 ± 0.10	1.4
	$56^\circ \lesssim b \lesssim 90^\circ$	0.14 ± 0.08	1.8

(a) Patches are centered in the indicated b coordinates.

(b) $\hat{\mathbf{a}}$ has units $\text{R} (\text{MJy/sr})^{-1}$.

(c) Our results are for the 15 GHz data.

quencies as well⁴.

Since we find substantially lower levels of free-free emission than Foreground X, the corresponding templates (100 μ m and WHAM) cannot be very highly correlated. We confirmed this by a direct cross-correlation analysis between these two templates, as seen though the Tenerife triple beam in the Tenerife observing region, and found the dimensionless correlation coefficient between the two maps to be in the range 5%–30% depending on Galactic latitude. The corresponding results in physical units (R/MJy sr⁻¹) are shown in Table 2. Since the statistical properties of these maps are not accurately known, we estimated the error bars by repeating the analysis with one of the templates replaced by $4 \times 36 = 144$ transformed maps, rotated around the Galactic axis by multiples of 10° and/or flipped vertically and/or horizontally. Significant correlations are found only for the $b > 20^\circ$ and $b > 30^\circ$ cuts, and the bulk of this correlation seems to come from a region of the sky between $30^\circ < b < 42^\circ$.

For comparison, see McCullough (1997), Kogut (1997) and dOC00 results shown in Table 2. This table indicates significant variations across the sky in the relationship between H_α and the 100 μ m emission, which could be related to variations in Hydrogen ionization fraction.

Finally, writing the frequency dependence as $a \propto \nu^\beta$ and recalling that the correlation coefficient is, by definition, $a = 1\text{K}/\mu\text{K} = 10^6$ for H82 at 408 MHz, we obtain the spectral index limits of $-2.9 \lesssim \beta \lesssim -3.6$ for the 10 GHz–H82 correlation and $-3.2 \lesssim \beta \lesssim -3.7$ for the 15 GHz–H82 correlation. These values are slightly steeper than the canonical sub-GHz slope of $-2.7 \lesssim \beta \lesssim -2.9$ (Davies *et al.* 1998; Platania *et al.* 1998), but consistent with a steepening of the spectrum of cosmic ray electrons at higher energies (Rybicki and Lightman 1979). The relatively high Tenerife synchrotron signal seen in Table 1 could be interpreted as slight spatial variability of the frequency dependence (Tegmark *et al.* 2000).

2.4. Power spectrum of foregrounds

Figure 1 shows the rms contribution of each foreground. In order to understand which angular scales contribute most to this rms, we compute the angular power spectra of the template emissions. The angular power spectra of the DIRBE, Haslam and WHAM components are shown in Figure 2 for the 20°–30° slice of Figure 1, scaled to 15 GHz using the correlation coefficients found above. All three power spectra are seen to be compatible with a power laws of ℓ^{-3} (solid line) at small angular scales ($\ell > 10 - 15$)⁵, while DIRBE and Haslam are seen to steepen further on very large scales ($\ell < 10 - 15$) to a power law closer to ℓ^{-6} (dashed line).

We remind the reader that there is, strictly speaking, no such thing as the power spectrum of a Galactic foreground, since the latitude dependence implies that it is not an isotropic random field. What we refer to here as the power spectrum of a foreground is simply the quantity that we care about in practice: its contribution to the




FIG. 2.— Angular power spectra for the DIRBE, Haslam and WHAM components as seen in the Tenerife data. Results are from the 20°–30° slice shown in Figure 1. The curves show ℓ^{-6} (dashed line) and ℓ^{-3} (solid line) power laws.

measurement of a CMB power spectrum. We have therefore computed the curves in Figure 2 by convolving the template maps with the Tenerife triple beam and treated the result as if it were a CMB map, computing quadratic band power estimators as described in Tegmark (1997) and Bond, Jaffe and Knox (1998).

3. AN ALTERNATIVE MODEL FOR FOREGROUND X

The results presented above show that free-free emission cannot explain Foreground X. Since there is presently only one other strong contender, spinning dust as proposed by DL98, the case would appear to be closed⁶. However, as we describe below, the situation seems to be a little more complicated.

100 μ m correlations have been detected at high significance on large ($3^\circ - 7^\circ$) angular scales: at 4.2σ at 31 GHz by DMR, at 6.6σ in the 19 GHz survey, at 6.4σ at 15 GHz by Tenerife and at 5.1σ at 10 GHz by Tenerife. These detections reveal a spectrum rising towards lower frequencies, with hints of a plateau or turnover between 10 and 15 GHz. On intermediate (degree) scales, the Saskatoon data gave a marginal detection while the QMAP data gave only an upper limit. At small (arcminute) scales the situation is even more confusing: although F01 recently reported a correlation of an HII region (LPH 201.663+1.643) and a dark cloud (L 1622) at 5, 8 and 10 GHz with the SFD98 100 μ map (Schlegel *et al.* 1998), these values are not in concordance with originally reported value given by OVRO at 14.5 GHz (Leitch *et al.* 1997)⁷. Moreover, only the dark cloud correlation detected by F01 seems to be clearly consistent with our (large scale) Tenerife result of $\hat{a} = 71.4 \pm 14.2 \mu\text{K}(\text{MJy}/\text{sr})^{-1}$ (see Table 1). This discrepancy cannot be blamed on variations in dust column density between different sky patches, since they would

⁴ A full DMR–WHAM correlation analysis is in progress (Kogut *et al.* 2002), so the results presented here should be viewed as preliminary. For all-sky cross-correlations, large scale variations in the WHAM map will need to be better accounted for.

⁵ Our DIRBE result is compatible with that reported by Wright (1998).

⁶ There is also the possibility of Foreground X is due to magnetic dipole emission from ferromagnetic grain materials (Draine and Lazarian 1999). This model can be ruled out if foreground X can be shown to correlate better with 12 μ m than 100 μ m emission.

⁷ The statistical significance of this result is unclear, since the OVRO measurement was published without error bars.




FIG. 3.— 12, 60 and 100 μ m IRAS images from the “mushroom” cloud MBM 20. These images are squares of $12.8^\circ \times 12.8^\circ$ centered at $l = 210.9, b = -34.5$. A simple visual comparison of these IRAS images at different frequencies suggests that although the large-scale features generally match up, small scale features can be quite different. Experiments such as ground-based interferometers, MAP and Planck LFI may have the angular resolution at the relevant low frequencies that are needed to be able to confirm or to rule out our grain segregation model.

not affect the dust-to-CMB conversion factor $\hat{\mathbf{a}}$. In summary, in average the most solid detections of Foreground X are all on scales of a few degrees or larger. Why is this foreground so elusive on smaller scales?

3.1. A grain size segregation model

Here we will argue in support of a model where Foreground X is spinning dust after all, and the problem is with the template used to find it, *i.e.*, 100 μ m dust emission.

Galactic dust grains come in a wide range of sizes (Weingartner and Draine 2000a). The 100 μ m template is dominated by large dust grains that are close to thermal equilibrium with the interstellar radiation field, and are radiating thermally at temperatures around 20K (Reach *et al.* 1995). In contrast, the small grains that can spin fast enough to produce the rotational emission of Draine and Lazarian cool faster than the mean time between two photon absorption events. Therefore, they spend most of their time near their ground state. They get heated to as much as a few hundred Kelvin when they absorb a photon, and radiate non-thermally with a spectrum that is both bluer and broader than for the large thermalized grains (Draine and Lee 1984). As a result, the emission from small grains peaks at shorter wavelengths like 10–30 μ m. Indeed, it is such non-equilibrium behavior that enables the Draine and Lazarian emission mechanism to work.

Previous work has generally assumed that the relative abundance of grain sizes is independent of position and that the 100 μ m map (dominated by large grains) is a good tracer for small grains as well. However, this seems to not be the case. A simple visual comparison of the IRAS maps at different frequencies suggests that although the large-scale features generally match up, small scale features can be quite different (see Figure 3).

Our proposed solution to the spinning dust puzzle is therefore that this component (small grains) only correlates well with 100 μ m emission (large grains) on fairly large angular scales. Occasional agreements on small scales would of course not be precluded (which could explain the F01 and Leitch *et al.* 1997 results), but should not be expected to hold in general.

As pointed out by Weingartner and Draine (2000b), it is not physically implausible for small and large dust grains to be separated. They showed that small dust grains can stay locked to the gas, while large dust grains in diffuse clouds can drift due to the effects of anisotropic starlight. After 10^7 yr, a typical lifetime of a diffuse interstellar cloud, such drifts can separate large and small grains by up to a few degrees at high Galactic latitudes.

3.2. Testing our model

A first prediction of this model is that shorter wavelength dust maps should trace spinning dust at least as well as a 100 μ m map (indeed, better on small scales)⁸. To test this, we repeated our analysis with the 100 μ m map replaced by DIRBE maps at other wavelengths. The results are shown in Figure 4 (lower curve). A first glance, it appears inconsistent with our prediction. However, contributions other than Galactic dust can clearly spoil the correlation, and several such contaminants are known to be present. Indeed, the figure shows that removing zodiacal emission from dust in our solar system increases the short wavelength correlations so much that the 60 μ m map traces spinning dust marginally better than the 100 μ m map (upper curve). The shorter wavelength maps also contain a substantial point source contribution. We used a new merged and destriped DIRBE+IRAS 12 μ m map⁹ to eliminate all 5- σ point source before convolving it with the Tenerife beam. Figure 4 shows that this new 12 μ m template traces spinning dust almost as well as the 100 μ m map (“Fink12 μ m”). For comparison, correlations with the 12 μ m map are shown in Table 1, bottom.

We tried two additional approaches to further increase the correlations (none of which helped more than marginally). The first was to apply a zodiacal cut on the zodi removed maps. The fact that this failed to increase

⁸ This prediction was also made by DL98, who argued that the 30 GHz emission and diffuse 12 μ m emission should be correlated, since both originate in grains containing ≈ 100 atoms.

⁹ This new map was constructed in the same fashion as SFD98 100 μ m map (for details about how such composite maps are made see Schlegel *et al.* (1998) or <http://www.astro.princeton.edu/~schlegel/dust/index.html>).




FIG. 4.— Dust correlations for the 12–240 μ m DIRBE maps at 15 GHz. Correlations of DIRBE maps with zodi are shown in blue, correlations of DIRBE maps with zodi removed are shown in red, and correlation with 12 μ m map with zodi and point source removed is shown in magenta.

correlations suggests that the zodi-removal performed by the DIRBE team was already adequate for our purposes. The second was to compute the linear combination of all DIRBE maps that gave the best correlation, to see if contaminants could be identified and subtracted spectrally. Correlations with this composite map are also shown in Table 1, bottom (see “Combo”). Our interpretation of these negative results is that our large dust grain templates are already tracing spinning dust quite well on large scales, so that it is impossible to do much better. Rather, the key tests will involve the correlation on small angular scales.

4. DISCUSSION

We have used the WHAM map to show that DIRBE-correlated microwave emission (Foreground X) cannot be explained as free-free emission. Since the spinning dust alternative has been challenged by recent small-scale observations, we have argued for an alternative model involving small scale grain segregation, where small and large dust grains are well correlated only on large angular scales.

We found that the zodi-cleaned 60 μ m DIRBE map traces Foreground X marginally better than the 100 μ m map, and that even the 12 μ m map does a good job after zodi and point source removal. Although these preliminary results are mildly encouraging, the smoking-gun test of this model will involve cross-correlating dust maps at various frequencies with high-resolution CMB data in the 10–30 GHz range.

Fortunately, such maps should be available shortly, from CMB experiments such as ground-based interferometers, the NASA MAP satellite¹⁰ and the Planck LFI¹¹. For instance, the expected sensitivity of the MAP 22 GHz channel is about 11 μ K per 56 arcminute pixel. For comparison, the expected small-scale fluctuations from spinning dust at 22 GHz are of order 20 μ K at high Galactic latitudes, ranging from a few to 10² μ K from a clean to a dirty re-

gion¹². Since MAP will measure tens of thousands of such pixels, it should be readily able to confirm or rule out our model. The prediction is that the small-scale signal will be substantially better traced by shorter wavelength dust maps. Since various contaminants may be important in these short wavelength maps (*e.g.*, polycyclic aromatic hydrocarbons – PAHs), it will also be worth performing a multicomponent fit using dust maps at all available frequencies, to find the linear combination of the dust maps that constitutes the best spinning dust template. High-resolution low frequency ground-based experiments such as CBI¹³ and DASI¹⁴ may also be able to test our hypothesis, but this is far from clear because they operate in Ka-band (26–36 GHz), where the spinning dust signal is small. In conclusion, we have proposed a resolution to the puzzle of Foreground X. Observations during the coming year should be able to test it.

We would like to thank Bruce Draine for useful comments. Support was provided by the NASA grant NAG5-9194 (*né* NAG5-6034) and the University of Pennsylvania Research Foundation. WHAM is supported by the NSF through grant AST96-19424. We acknowledge the NASA office of Space Sciences, the COBE flight team, and all those who helped process and analyze the DIRBE data.

REFERENCES

- Bennett, C.L., et al. 1992, ApJ, 396, L7
- Boggess, N.W., et al. 1992, ApJ, 397, 420
- Bond, J.R., Jaffe, A.H., Knox, L.E. 1998, Phys. Rev. D, 57, 2117
- Davies R.D., Watson, R.A., and Gutierrez, C.M. 1998, MNRAS, 278, 925
- de Oliveira-Costa, A., et al. 1997, ApJ, 482, L17
- de Oliveira-Costa, A., et al. 1998, ApJ, 509, L9
- de Oliveira-Costa, A., et al. 1999, ApJ, 527, L9 (dOC99)
- de Oliveira-Costa, A., et al. 2000, ApJ, 542, L5 (dOC00)
- Draine, B.T., and Lazarian, A. 1998, ApJ, 494, L19 (DL98)
- Draine, B.T., and Lazarian, A. 1999, ApJ, 512, 740
- Draine, B.T., and Lee, H.M. 1984, ApJ, 285, 89
- Finkbeiner, D.P., Davis, M., Frank, C., Schlegel, D.J., and Heiles, C. 2001, astro-ph/0109534, ApJL in press (F01)
- Gutiérrez, C.M., et al. 2000, ApJ, 529, 47
- Haffner, L.M. et al. 1999, ApJ, 523, 223
- Haslam, C.G.T., et al. 1982, A&AS 47, 1 (H82)
- Kogut, A. 1997, AJ, 114, 1127
- Kogut, A., et al. 1996a, ApJ, 460, 1
- Kogut, A., et al. 1996b, ApJ, 464, L5
- Kogut, A., et al. 2002, in preparation
- Leitch, E.M., et al. 1997, ApJ, 486, L23
- McCullough, P.R. 1997, AJ, 113, 2186
- Mukherjee, P. et al. 2000, astro-ph/0002305
- Platanica, P., et al. 1998, ApJ, 505, 473
- Reach, W.T. et al. 1995, ApJ, 451, 188
- Reich, W. 1982, A&AS, 48, 219
- Reich, P., and Reich, W. 1986, A&AS, 63, 205
- Rybicki, G. B. and Lightman, A. P. 1979, Radiative Processes in Astrophysics (New York: Wiley), p.174
- Schlegel, D.J., Finkbeiner, D.P., and Davis, M. 1998, ApJ, 500, 525 (SFD98)
- Tegmark, M. 1997, Phys. Rev. D, 56, 4514
- Tegmark, M. et al. 2000, ApJ, 530, 133
- Weingartner, J.C., and Draine, B.T. 2000a, astro-ph/0008146
- Weingartner, J.C., and Draine, B.T. 2000b, astro-ph/0010117
- Wright, E.L. 1998, ApJ, 496, 1

¹²We made this estimate by high-pass filtering the Fink12 μ m map to the degree scale – this was done by taking the difference of two Fink12 μ m maps, one smoothed by 0.7° and another by 2°. Using $\tilde{a} \approx 760\mu\text{m}/(\text{MJy/sr})$ from Table 1, we converted this difference map to μK , and finally made a small spectral correction from 15 to 22 GHz.

¹³<http://astro.caltech.edu/~tjp/CBI/>

¹⁴<http://astro.uchicago.edu/dasi/>

¹⁰<http://map.gsfc.nasa.gov/>

¹¹<http://astro.estec.esa.nl/SA-general/Projects/Planck/>